



ELSEVIER

# Nonsense-mediated mRNA decay: terminating erroneous gene expression

Kristian E Baker and Roy Parker<sup>1</sup>

Nonsense-mediated mRNA decay is a surveillance pathway that reduces errors in gene expression by eliminating aberrant mRNAs that encode incomplete polypeptides. Recent experiments suggest a working model whereby premature and normal translation termination events are distinct as a consequence of the spatial relationship between the termination codon and mRNA binding proteins, a relationship partially established by nuclear pre-mRNA processing. Aberrant termination then leads to both translational repression and an increased susceptibility of the mRNA to multiple ribonucleases.

## Addresses

Howard Hughes Medical Institute, University of Arizona, 1007 East Lowell Street, Tucson, Arizona 85721 USA

<sup>1</sup>e-mail: rrparker@u.arizona.edu

**Current Opinion in Cell Biology** 2004, **16**:293–299

This review comes from a themed issue on Nucleus and gene expression  
Edited by Elisa Izaurralde and David Spector

0955-0674/\$ – see front matter  
© 2004 Elsevier Ltd. All rights reserved.

DOI 10.1016/j.j.ceb.2004.03.003

## Abbreviations

<b>CBP</b>	cap binding protein
<b>DSE</b>	downstream sequence element
<b>EJC</b>	exon–exon junction complex
<b>mRNP</b>	messenger ribonucleoprotein particle
<b>NMD</b>	nonsense-mediated mRNA decay
<b>ORF</b>	open reading frame
<b>PARN</b>	poly(A) ribonuclease
<b>PTC</b>	premature translation termination codon
<b>RNP</b>	ribonucleoprotein
<b>UTR</b>	untranslated region

## Introduction

Eukaryotic cells exhibit quality-control mechanisms that recognize and degrade mRNAs that have not completed nuclear pre-mRNA processing or that fail to encode a polypeptide faithful to the gene sequence. Such aberrant mRNAs are degraded rapidly, presumably before the accumulation of deviant protein products that could have adverse effects on the cell. One such quality-control system is referred to as nonsense-mediated mRNA decay (NMD). Although the first mRNA substrates identified for NMD encoded premature translation termination codons (PTCs; see for example [1,2]), it is now appreciated that NMD degrades a variety of aberrant mRNAs that arise as a result of either mutations or defects in pre-

mRNA processing. The common theme is that substrates of NMD manifest alterations in the normal spatial relationship between the termination codon and additional RNA features. For example, NMD substrates include transcripts with an upstream ORF in the 5' UTR [3], extended 3' UTRs (which alters the spatial relationship between the termination codon and the 3' poly(A) tail [4–7]), bicistronic mRNAs [8\*\*], and mRNAs with additional, and utilized, out-of-frame AUG codons, which lead to premature translation termination as a consequence of translational recoding or leaky ribosome scanning [9].


NMD can target a substantial part of the eukaryotic transcriptome and can be biologically important. For example, microarray experiments suggest that the levels of ~10% of yeast transcripts are affected by alterations in NMD [8\*\*,10]. Furthermore, analysis of mammalian splicing suggests that possibly one-third of alternative splicing events produce transcripts that are substrates for NMD [11\*,12]. The importance of mRNA surveillance is exemplified by the expression of truncated polypeptides that result in gain-of-function or dominant-negative mutant proteins having deleterious effects [13–15]. Therefore, it is consequential to understand how events surrounding translation termination contribute to the recognition of normal versus aberrant transcripts and how recognition is communicated into rapid degradation of the mRNA. In this review, we will present recent developments which provide insight into these molecular events that contribute to NMD.

## Translational termination: distinct events for normal versus aberrant mRNAs

A critical distinction between premature, or aberrant, and normal translation termination appears to be a biochemical difference in the termination reaction itself; this difference then elicits downstream consequences for the fate of the mRNA. The possibility that termination events could be distinct was first suggested from evidence that a ribosome positioned at a termination codon can terminate, read through, or resume scanning after termination, depending on the termination context [16,17].

Evidence that NMD involves alterations in the termination process has come from the analysis of proteins that are essential for NMD, which include Upf1, Upf2 and Upf3 [18]. Upf1 protein has RNA-dependent ATPase and 5'–3' helicase activities which are required for NMD and presumably promote an unknown RNA–RNA or ribonucleoprotein (RNP) rearrangement. Importantly, Upf

Figure 1



Models for nonsense-mediated decay of mRNA. Nuclear pre-mRNA processing events fashion mRNPs that are exported to the cytoplasm for translation. During the pioneer round of translation, ribosomes displace RNPs from the entire length of the coding region for wild-type mRNAs (left panel). The absence of certain RNPs and/or positive signals from remaining RNPs command proper translational termination, continued

proteins interact directly with the translation termination factors eRF1 and/or eRF3, providing a direct link between the termination complex and the NMD machinery [19,20]. Moreover, this interaction appears to be functionally important: *upf1Δ*, *upf2Δ* or *upf3Δ* strains show an increased frequency of nonsense suppression (i.e. translational readthrough) at termination codons in a variety of yeast genes [21,22]. These data suggest that Upf proteins cycle between a free state and an interaction with terminating ribosomes. Interestingly, in *C. elegans*, *Drosophila* and mammals, additional proteins are required for NMD [23]. These proteins are involved in catalyzing a cycle of Upf1 phosphorylation and dephosphorylation required for NMD [24,25,26<sup>\*</sup>,27<sup>\*</sup>,28<sup>\*\*</sup>,29<sup>\*</sup>]. It is proposed that brief phosphorylation of Upf1 occurs during translational termination, as PP2a, the Upf1 phosphatase, interacts with eRF1 [30]. In addition, Upf1 is hyperphosphorylated when associated with polysomes [31]. The phosphorylation cycle may modulate the ATPase or helicase activity of Upf1 and/or the remodeling of the ribosome-associated termination event [29<sup>\*</sup>].

Compelling evidence that premature and normal translation termination events are distinct has come from a biochemical analysis of termination. A critical observation is that the ribosome toeprint generated at a normal translation termination codon is distinct from the toeprint of a ribosome terminating at a premature termination codon (Allan Jacobson, personal communication); moreover, this distinction is dependent upon the presence of Upf1p at the premature termination codon. These data suggest that premature translation termination events in the presence of Upf1p will be aberrant, whereas 'normal' termination proceeds in the absence of Upf function. Given this, an important question is what features of a messenger ribonucleoprotein particle (mRNP) define a translation termination event as premature versus normal.

### Translation termination and organization of the mRNP

A critical determinant of whether an mRNA is subjected to NMD is the relationship between the translation termination codon and other mRNP features. In mammalian mRNAs, stop codons at distances greater than 50–55 nucleotides upstream from the 3'-most exon-exon junction are perceived as premature and lead to NMD [32–35]. The coupling between RNA splicing and NMD is dictated by the splicing-dependent deposition of a protein complex, referred to as the exon-exon junction complex (EJC), ~20–24 nucleotides upstream

of the 5' splice site following splicing [36–40]. Furthermore, an interaction between the EJC protein Y14 and Upf3 is required to trigger mRNA decay, thus providing a physical and functional link between the EJC and NMD [41<sup>\*\*</sup>].

An mRNA can also be recognized as aberrant through mRNP features independent of an EJC, as is the case for several mammalian genes [42]. Moreover, in yeast and *Drosophila*, for example, the recognition of a PTC is independent of an intron and RNA splicing [28<sup>\*\*</sup>,43,44]. In yeast, evidence suggests that specific sequences within the coding region, referred to as downstream sequence elements (DSEs), are required to trigger NMD when found 3' of a translation termination codon [45]. Such elements have been proposed to bind Hrp1p, which in turn interacts with Upf1p, and which, like the EJC complex, may recruit additional NMD factors to the mRNA and thereby trigger decay [46]. However, there may be additional features of yeast mRNAs that distinguish normal and aberrant transcripts, as a nonsense codon downstream of a well-characterized DSE can still trigger NMD [47<sup>\*</sup>]. An unresolved issue is whether the various mRNP features influence the nature of termination, or function to trigger NMD in a step following termination.

A popular model for NMD suggests that the first ribosome to translate the mRNA removes the EJC or other mRNP complexes from along the entire coding region, and that if such 'coding region' complexes remain on the mRNA following termination, NMD is activated. This is supported by the observation that translating ribosomes can remove the EJC component Y14 [48]. Evidence that EJC remodeling occurs during the first, or pioneer, round of translation includes observations that NMD reduces a pool of mRNAs bound to CBP80 and CBP20, components of the nuclear cap-binding complex [49,50]. Recently, eIF4AIII, a homologue of the translational initiation factor eIF4AI, was identified as a component for NMD in mammalian cells [51<sup>\*</sup>,52]. Additionally, eIF4AIII is primarily a nuclear protein and interacts with eIF4G and eIF4B. It remains to be established, however, whether eIF4AIII provides a link between the first round of mRNA translation and NMD, or whether its role is more indirect.

Two divergent views have been put forth to explain how protein complexes remaining on the mRNP might trigger NMD (Figure 1). In one, ribosomes terminating at pre-

---

**(Figure 1 Legend Continued)** translation and mRNA stability. Failure of the ribosome to remove an RNP as a consequence of a PTC (right panel) may lead to either appropriate translational termination and 'Surveillance complex' scanning of the mRNA, or improper termination as a result of negative signals from the downstream RNP combined with the failure of other positive signals to be communicated to the ribosome. Interactions between an inappropriate RNP and the 'surveillance complex' or ribosome result in translational repression and rapid degradation involving multiple decay pathways.

mature stop codons deposit a Upf-containing protein complex, referred to as the 'Surveillance complex', on the mRNA; this protein complex scans 3' for EJCs or DSEs and if one is encountered, NMD is triggered [53]. An alternative model suggests that the mRNP features mark the context of the normal termination codon in a manner that leads to proper termination and mRNA stabilization [17]. In the absence of such positive interactions, termination is aberrant and the mRNA is targeted for NMD.

Several observations suggest that the poly(A) tail and its associated binding protein, Pab1p, may constitute a positive factor that affects the nature of termination and the process of NMD. First, in yeast, the poly(A) tail is generally proximal to the termination codon and could provide contextual information. In addition, Pab1p interacts with eRF3, can influence the nature of translation termination, and may even promote ribosome recycling [54]. Similarly, in mammals the poly(A) tail might interact with the last EJC as a result of exon definition and thereby also provide an important mark that enhances the function of the mRNA. Consistent with the EJC and Upf proteins having positive roles in mRNA function, the presence of either an EJC or a Upf protein within the coding region serves to enhance mRNA 3'-end formation and increases translation yield of the mRNA [55,56,57\*\*]. These data hint at the possibility that NMD might arise, at least in part, from a failure of the terminating ribosome to interact with Pab1p.

### Recognition of aberrant mRNAs and downstream events

The recognition of an mRNA as aberrant by the NMD machinery leads to a series of changes in the behavior of the mRNA. Primarily, being recognized as nonsense-containing increases the susceptibility of an mRNA to multiple mRNA degradation systems. Two general degradation pathways for normal mRNAs have been identified in eukaryotic cells [58]. In both, decay of the mRNA begins with shortening of the 3' poly(A) tail followed by 3'-to-5' degradation and/or decapping and 5'-to-3' degradation. In yeast, NMD triggers rapid deadenylation, removes the inhibition of decapping by the poly(A) tail, and accelerates the rate of decapping [47\*,59]. These events result in the mRNA being degraded primarily by rapid deadenylation-independent decapping. However, when decapping and 5'-to-3' decay is blocked, NMD promotes faster decay of mRNAs through the 3'-to-5' decay pathway [47\*,59,60\*,61\*]. Kinetic analysis provides evidence that NMD accelerates 3'-to-5' decay primarily by a change in deadenylation rates, although there may be a small effect on the rate of 3'-to-5' decay of the mRNA body [47\*,61\*].

The decapping and 3'-to-5' decay pathways in mammals are also stimulated by NMD. For example, mammalian

NMD increases the rate of deadenylation and decapping [62,63\*\*,64]. In addition, depletion of the decapping enzyme, Dcp2, or of Rrp6p or PARN, proteins putatively required for 3'-to-5' degradation of mRNAs, leads to an increase in the abundance of steady-state nonsense-containing mRNAs [63\*]. With evidence that both the 5'-to-3' and 3'-to-5' degradation pathways can act separately on NMD substrates in yeast or mammals, the question arises of how defects in either pathway can cause a decrease in the decay of NMD substrates.

NMD can also target an mRNA for enhanced rates of endonucleolytic cleavage. For example, in erythrocytes, NMD can trigger faster susceptibility of the  $\beta$ -globin mRNA to cleavage by the PMR1-like endonuclease [65,66]. Thus, even though there exists conservation of the proteins required for NMD, there are several different mechanisms by which NMD accelerates RNA degradation.

The existence of multiple degradation pathways for NMD substrates indicates that NMD leads to dramatic changes in the status of the mRNA. An appealing possibility is that NMD leads to targeting of the mRNA to a non-translating pool of transcripts. Because translation factors and translation itself appear to protect mRNAs from destruction, this would generally increase the instability of the transcript. Consistent with this view, measuring the translation efficiency of nonsense-containing mRNAs demonstrated that Upf1p plays a role in repressing the translation of the mRNA, independent of mRNA degradation [7]. This argues that one consequence of NMD is to render the RNA translationally inactive and thereby susceptible to the various decay pathways.

### Physical interactions between the NMD, translation and decay machineries

An emerging concept is that NMD leads to both translational repression and the recruitment of ribonucleases. The reduction in translation could be due to aberrant translation termination *per se*, or may be a more direct result of the function of the Upf proteins. For example, proper termination may enhance the efficiency of subsequent rounds of translation by promoting ribosome recycling to the initiation complex, whereas aberrant termination would fail to complete this function and lead to reduced translation. By contrast, the Upf2 protein has been found to interact physically with eIF4A and eIF1 [43]. Since mutations in the yeast eIF1 protein inhibit NMD [67], this interaction may be part of an active mechanism by which aberrant mRNAs are translationally repressed.

Several observations also provide evidence for a more direct interaction between the Upf proteins and the mRNA degradation machinery. For example, yeast Upf1p shows a strong two-hybrid interaction with Dcp2p,

a subunit of the decapping enzyme [68]. Similarly, mammalian decapping enzyme co-immunoprecipitates with Upf proteins [63<sup>••</sup>,69]. Biochemical interactions have also been detected between Upf1 and factors involved in 3'-to-5' degradation of mRNAs. In mammalian cells, Upf1 can co-immunoprecipitate with components of the exosome, which is the nuclease complex responsible for 3'-to-5' degradation of mRNAs [63<sup>••</sup>]. In addition, there exists a physical interaction between yeast Upf1p and Ski7p, an exosome-associated protein that may adapt the nuclease complex to different substrates [61<sup>•</sup>]. These physical interactions suggest that Upf proteins may play an additional, and more direct, role in recruiting nucleases to the mRNA.

### Conclusions and future directions

Over the last few years there has been significant progress in understanding substrate recognition and in characterizing the protein players required for NMD. However, an increased comprehension of the molecular interactions between the Upf proteins, translational termination and initiation factors, and the decay machinery is needed to reveal the actual molecular mechanism of NMD. In addition, there are several broader issues that still require consideration. For example, where does NMD occur in the cell and what constraints does the location put on the mechanism? In mammalian cells, NMD can affect both nuclear and cytoplasmic mRNA levels, suggesting either that translation and NMD can occur in the nucleus, or that there is feedback following cytoplasmic NMD. In yeast, and possibly mammals, discrete cytoplasmic foci, referred to as P-bodies, have been identified, in which normal decapping and 5'-to-3' degradation can occur [70<sup>•</sup>,71–73]. Does NMD also occur in P-bodies, and if so, what regulates formation or mRNA entrance into P-bodies? Similarly, it remains unclear how, in mammalian cells, the recognition of an mRNA as nonsense-containing can affect events in the nucleus including splicing, abundance of the nuclear mRNA pool, and retention of transcripts at the gene [74]. Understanding these issues should not only lead to insights into the process of NMD, but also reveal general properties of mRNA biology in eukaryotic cells.

### Acknowledgements

We would like to thank Elisa Izaurralde, Lynne Maquat, Allan Jacobson, Melissa Moore, Thierry Grange, David Bedwell, Gideon Dreyfuss and Nahum Sonenberg for communicating data and correspondence before publication. This work was supported by the Howard Hughes Medical Institute.

### References and recommended reading

Papers of particular interest, published within the annual period of review, have been highlighted as:

- of special interest
  - of outstanding interest
1. Losson R, Lacroute F: **Interference of nonsense mutations with eukaryotic messenger mRNA stability.** *Proc Natl Acad Sci U S A* 1979, **76**:5134-5137.
  2. Maquat LE, Kinniburgh AJ, Rachmilewitz EA, Ross J: **Unstable  $\beta$ -globin mRNA in mRNA-deficient  $\beta^0$ -thalassemia.** *Cell* 1981, **27**:543-553.
  3. Oliveira CC, McCarthy JE: **The relationship between eukaryotic translation and mRNA stability. A short upstream open reading frame strongly inhibits translational initiation and greatly accelerates mRNA degradation in the yeast *Saccharomyces cerevisiae*.** *J Biol Chem* 1995, **270**:8936-8943.
  4. Zaret KS, Sherman F: **Mutationally altered 3' ends of yeast *CYC1* mRNA affect transcript stability and translational efficiency.** *J Mol Biol* 1984, **177**:107-135.
  5. Pulak RA, Anderson P: **Structures of spontaneous deletions in *Caenorhabditis elegans*.** *Mol Cell Biol* 1988, **8**:3748-3754.
  6. Pulak R, Anderson P: **mRNA surveillance by the *Caenorhabditis elegans smg* genes.** *Genes Dev* 1993, **7**:1885-1897.
  7. Muhlrud D, Parker R: **Aberrant mRNAs with extended 3' UTRs are substrates for rapid degradation by mRNA surveillance.** *RNA* 1999, **5**:1299-1307.
  8. He F, Li X, Spatrick P, Casillo R, Dong S, Jacobson A: **Genome-wide analysis of mRNAs regulated by the nonsense-mediated and 5' to 3' mRNA decay pathways in yeast.** *Mol Cell* 2003, **12**:1439-1452.
  - The complement of transcripts regulated by NMD in yeast is identified and determined to be degraded primarily by the 5'-to-3' decay pathway.
  9. Welch EM, Jacobson A: **An internal open reading frame triggers nonsense-mediated decay of the yeast SPT mRNA.** *EMBO J* 1999, **18**:6134-6145.
  10. Lelivelt MJ, Culbertson MR: **Yeast Upf proteins required for RNA surveillance affect global expression of the yeast transcriptome.** *Mol Cell Biol* 1999, **19**:6710-6719.
  11. Lewis BP, Green RE, Brenner SE: **Evidence for the widespread coupling of alternative splicing and nonsense-mediated mRNA decay in humans.** *Proc Natl Acad Sci U S A* 2003, **100**:189-192.
  - One-third of alternatively spliced mRNAs in mammalian cells contain PTCs and are substrates for NMD.
  12. Green RE, Lewis BP, Hillman RT, Blanchette M, Lareau LF, Garnett AT, Rio DC, Brenner SE: **Widespread predicted nonsense-mediated mRNA decay of alternatively spliced transcripts of human normal and disease genes.** *Bioinformatics* 2003, **19**:1118-1121.
  13. Culbertson MR: **RNA surveillance. Unforeseen consequences for gene expression, inherited genetic disorders and cancer.** *Trends Genet* 1999, **15**:74-80.
  14. Hentze MW, Kulozik AE: **A perfect message: RNA surveillance and nonsense-mediated decay.** *Cell* 1999, **96**:307-310.
  15. Frischmeyer PA, Dietz HC: **Nonsense-mediated mRNA decay in health and disease.** *Hum Mol Genet* 1999, **8**:1893-1900.
  16. Janzen DM, Geballe AP: **Modulation of translation termination mechanisms by cis- and trans-acting factors.** *Cold Spring Harb Symp Quant Biol* 2001, **66**:459-467.
  17. Hilleren P, Parker R: **Mechanisms of mRNA surveillance in eukaryotes.** *Annu Rev Genet* 1999, **33**:229-260.
  18. Culbertson MR, Leeds PF: **Looking at mRNA decay pathways through the window of molecular evolution.** *Curr Opin Genet Dev* 2003, **13**:207-214.
  19. Czaplinski K, Ruiz-Echevarria MJ, Paushkin SV, Han X, Weng Y, Perlick HA, Dietz HC, Ter-Avanesyan MD, Peltz SW: **The surveillance complex interacts with the translation release factors to enhance termination and degrade aberrant mRNAs.** *Genes Dev* 1998, **12**:1665-1677.
  20. Wang W, Czaplinski K, Rao Y, Peltz SW: **The role of Upf proteins in modulating the translation read-through of nonsense-containing transcripts.** *EMBO J* 2001, **15**:880-890.
  21. Bidou L, Stahl G, Hatin I, Namy O, Rousset JP, Farabaugh PJ: **Nonsense-mediated decay mutants do not affect programmed-1 frameshifting.** *RNA* 2000, **6**:952-961.

22. Keeling KM, Lanier J, Du M, Salas-Marco J, Gao L, Kaenjak-Angeletti A, Bedwell DM: **Leaky termination at premature stop codons antagonizes nonsense-mediated mRNA decay in *S. cerevisiae*.** *RNA* 2004, in press.
23. Maquat LE: **Nonsense-mediated mRNA decay: a comparative analysis of different species.** *Curr Genomics* 2004, in press.
24. Page MF, Carr B, Anders KR, Grimson A, Anderson P: **SMG-2 is a phosphorylated protein required for mRNA surveillance in *Caenorhabditis elegans* and related to Upf1p of yeast.** *Mol Cell Biol* 1999, **19**:5943-5951.
25. Yamashita A, Ohnishi T, Kashima I, Taya Y, Ohno S: **Human SMG-1, a novel phosphatidylinositol 3-kinase-related protein kinase, associates with components of the mRNA surveillance complex and is involved in the regulation of nonsense-mediated mRNA decay.** *Genes Dev* 2001, **15**:2215-2228.
26. Chiu SY, Serin G, Ohara O, Maquat LE: **Characterization of human Smg5/7a: a protein with similarities to *Caenorhabditis elegans* SMG5 and SMG7 that functions in the dephosphorylation of Upf1.** *RNA* 2003, **9**:77-87.
- Human SMG6 functions in the dephosphorylation of hUpf1 and copurifies with the catalytic subunit of protein phosphatase 2A.
27. Anders KR, Grimson A, Anderson P: **SMG-5, required for *C. elegans* nonsense-mediated mRNA decay, associates with SMG-2 and protein phosphatase 2A.** *EMBO J* 2003, **22**:641-650.
- SMG-5 is required for NMD in *C. elegans* and serves to direct protein phosphatase 2A to SMG-2/UPF1 for dephosphorylation.
28. Gatfield D, Unterholzner L, Ciccarelli FD, Bork P, Izaurralde E: **Nonsense-mediated mRNA decay in *Drosophila*: at the intersection of the yeast and mammalian pathways.** *EMBO J* 2003, **22**:3960-3970.
- In *Drosophila*, the UPF1, UPF2, UPF3, SMG1, SMG5 and SMG6 proteins are required for NMD, and PTC definition occurs independently of exon boundaries and components of the EJC.
29. Ohnishi T, Yamashita A, Kashima I, Schell T, Anders KR, Grimson A, Hachiya T, Hentze MW, Anderson P, Ohno S: **Phosphorylation of hUPF1 induces formation of mRNA surveillance complexes containing hSMG-5 and hSMG-7.** *Mol Cell* 2003, **12**:1187-1200.
- Human UPF1 forms a complex with hSMG-5, hSMG-7 and protein phosphatase 2A that results in hUPF1 dephosphorylation, an event required for NMD.
30. Andjelkovic N, Zolnierowicz S, Van Hoof C, Goris J, Hemmings BA: **The catalytic subunit of protein phosphatase 2A associates with the translation termination factor eRF1.** *EMBO J* 1996, **15**:7156-7167.
31. Pal M, Ishigaki Y, Nagy E, Maquat LE: **Evidence that phosphorylation of human Upf1 protein varies with intracellular location and is mediated by a wortmannin-sensitive and rapamycin-sensitive PI3-kinase-related kinase signaling pathway.** *RNA* 2001, **7**:5-15.
32. Cheng J, Belgrader P, Zhou X, Maquat LE: **Introns are cis effectors of the nonsense-codon-mediated reduction in nuclear mRNA abundance.** *Mol Cell Biol* 1994, **14**:6317-6325.
33. Zhang J, Sun X, Qian Y, Maquat LE: **Intron function in the nonsense-mediated decay of  $\beta$ -globin mRNA: indications that pre-mRNA splicing in the nucleus can influence mRNA translation in the cytoplasm.** *RNA* 1998, **4**:801-815.
34. Zhang J, Sun X, Qian Y, LaDuca JP, Maquat LE: **At least one intron is required for the nonsense-mediated decay of triosephosphate isomerase mRNA: a possible link between nuclear splicing and cytoplasmic translation.** *Mol Cell Biol* 1998, **18**:5272-5283.
35. Thermann R, Neu-Yilik G, Deters A, Frede U, Wehr K, Hagemeier C, Hentze MW, Kulozik AE: **Binary specification of nonsense codons by splicing and cytoplasmic translation.** *EMBO J* 1998, **17**:3484-3494.
36. Le Hir H, Moore MJ, Maquat LE: **Pre-mRNA splicing alters mRNP composition: evidence for stable association of proteins at exon-exon junctions.** *Genes Dev* 2000, **14**:1098-1108.
37. Le Hir H, Izaurralde E, Maquat LE, Moore MJ: **The spliceosome deposits multiple proteins 20-24 nucleotides upstream of mRNA exon-exon junctions.** *EMBO J* 2000, **19**:6860-6869.
38. Kataoka N, Yong J, Kim VN, Velazquez F, Perkinson RA, Wang F, Dreyfuss G: **Pre-mRNA splicing imprints mRNA in the nucleus with a novel RNA-binding protein that persists in the cytoplasm.** *Mol Cell* 2000, **6**:673-682.
39. Lykke-Andersen J, Shu MD, Steitz JA: **Human Upf proteins target an mRNA for nonsense-mediated decay when bound downstream of a termination codon.** *Cell* 2000, **103**:1121-1131.
40. Lykke-Andersen J, Shu MD, Steitz JA: **Communication of the position of exon-exon junctions to the mRNA surveillance machinery by the protein RNPS1.** *Science* 2001, **293**:1836-1839.
41. Gehring NH, Neu-Yilik G, Schell T, Hentze MW, Kulozik AE: **Y14 and hUPF3b form an NMD-activating complex.** *Mol Cell* 2003, **11**:939-949.
- An interaction between hUPF3b and the EJC protein Y14 is essential for NMD in mammalian cells.
42. Maquat LE: **Nonsense-mediated mRNA decay: splicing, translation and mRNP dynamics.** *Nat Rev* 2004, **5**:89-99.
43. Mendell JT, Medghalchi SM, Lake RG, Noensie EN, Dietz HC: **Novel Upf2p orthologues suggest a functional link between translation initiation and nonsense surveillance complexes.** *Mol Cell Biol* 2000, **20**:8944-8957.
44. Gonzalez CI, Bhattacharya A, Wang W, Peltz SW: **Nonsense-mediated mRNA decay in *Saccharomyces cerevisiae*.** *Gene* 2001, **274**:15-25.
45. Peltz SW, Brown AH, Jacobson A: **mRNA destabilization triggered by premature translational termination depends on at least three cis-acting sequence elements and one trans-acting factor.** *Genes Dev* 1993, **7**:1737-1754.
46. Gonzalez CI, Ruiz-Echevarria MJ, Vasudevan S, Henry MF, Peltz SW: **The yeast hnRNP-like protein Hrp1/Nab4 marks a transcript for nonsense-mediated mRNA decay.** *Mol Cell* 2000, **5**:489-499.
47. Cao D, Parker R: **Computational modeling and experimental analysis of nonsense-mediated decay in yeast.** *Cell* 2003, **113**:533-545.
- Nonsense-containing mRNAs are targeted for deadenylation-independent decapping, accelerated deadenylation and faster rates of decapping in yeast.
48. Dostie J, Dreyfuss G: **Translation is required to remove Y14 from mRNAs in the cytoplasm.** *Curr Biol* 2002, **12**:1060-1067.
49. Ishigaki Y, Li X, Serin G, Maquat LE: **Evidence for a pioneer round of mRNA translation: mRNAs subject to nonsense-mediated decay in mammalian cells are bound by CBP80 and CBP20.** *Cell* 2001, **106**:607-617.
50. Lejeune F, Ishigaki Y, Li X, Maquat LE: **The exon junction complex is detected on CBP80-bound but not eIF4E-bound mRNA in mammalian cells: dynamics of mRNP remodeling.** *EMBO J* 2002, **21**:3536-3545.
51. Palacios IM, Gatfield D, St Johnston D, Izaurralde E: **An eIF4AIII-containing complex required for mRNA localization and nonsense-mediated mRNA decay.** *Nature* 2004, **427**:753-757.
- The homologue of translation initiation factor eIF4AI, eIF4AIII, is a component of the EJC and is required for NMD in mammalian cells.
52. Chan CC, Dostie J, Diem MD, Feng W, Mann M, Rappsilber J, Dreyfuss G: **eIF4A3 is a novel component of the exon junction complex.** *RNA* 2004, **10**:200-209.
53. Gonzalez CI, Bhattacharya A, Wang W, Peltz SW: **Nonsense-mediated mRNA decay in *Saccharomyces cerevisiae*.** *Gene* 2001, **274**:15-25.
54. Mangus DA, Evans MC, Jacobson A: **Poly(A)-binding proteins: multifunctional scaffolds for the post-transcriptional control of gene expression.** *Genome Biol* 2003, **4**:223.
55. Matsumoto K, Wassarman KM, Wolfe AP: **Nuclear history of a pre-mRNA determines the translational activity of cytoplasmic mRNA.** *EMBO J* 1998, **17**:2107-2121.
56. Nott A, Meislin SH, Moore MJ: **A quantitative analysis of intron effects on mammalian gene expression.** *RNA* 2003, **9**:607-617.

57. Nott A, Le Hir H, Moore MJ: **Splicing enhances translation in mammalian cells: an additional function of the exon junction complex.** *Genes Dev* 2004, **18**:210-222.  
EJC protein deposition within an mRNA ORF promotes polysome association and an increased translational yield.
58. Parker R, Song H: **The enzymes and control of eukaryotic mRNA turnover.** *Nat Struct Mol Biol* 2004, **11**:121-127.
59. Muhrad D, Parker R: **Premature translational termination triggers mRNA decapping.** *Nature* 1994, **370**:578-581.
60. Mitchell P, Tollervey D: **An NMD pathway in yeast involving accelerated deadenylation and exosome-mediated 3'-5' degradation.** *Mol Cell* 2003, **11**:1405-1413.  
NMD can target mRNAs to deadenylation and exosome-mediated exonucleolytic decay in a Upf1- and translation-dependent manner.
61. Takahashi S, Araki Y, Sakuno T, Katada T: **Interaction between Ski7p and Upf1p is required for nonsense-mediated 3'-to-5' mRNA decay in yeast.** *EMBO J* 2003, **22**:3951-3959.  
NMD substrates are targeted to accelerated 3'-to-5' decay catalyzed by the exosome. Upf1p interaction with exosome-associated protein Ski7p is functionally important for 3'-to-5' decay.
62. Chen CY, Shyu AB: **Rapid deadenylation triggered by a nonsense codon precedes decay of the RNA body in a mammalian cytoplasmic nonsense-mediated decay pathway.** *Mol Cell Biol* 2003, **23**:4805-4813.
63. Lejeune F, Li X, Maquat LE: **Nonsense-mediated mRNA decay in mammalian cells involves decapping, deadenylating, and exonucleolytic activities.** *Mol Cell* 2003, **12**:675-687.  
Nonsense-containing mRNAs are stabilized in mammalian cells depleted for the decapping enzyme Dcp2 or for PM/Sci100 and PARN, proteins involved in 3'-to-5' RNA degradation. Human Upf proteins coimmunoprecipitate with the decapping enzyme, the 5'-to-3' exonuclease Xrn1, PARN, and several exosomal proteins.
64. Couttet P, Grange T: **Premature termination codons enhance mRNA decapping in human cells.** *Nucleic Acids Res* 2004, **32**:488-494.
65. Stevens A, Wang Y, Bremer K, Zhang J, Hoepfner R, Antoniou M, Schoenberg DR, Maquat LE:  **$\beta$ -globin mRNA decay in erythroid cells: UG site-preferred endonucleolytic cleavage that is augmented by a premature termination codon.** *Proc Natl Acad Sci U S A* 2002, **99**:12741-12746.
66. Bremer KA, Stevens A, Schoenberg DR: **An endonuclease activity similar to *Zenopus* PMR1 catalyzes the degradation of normal and nonsense-containing human  $\beta$ -globin mRNA in erythroid cells.** *RNA* 2003, **9**:1157-1167.
67. Cui Y, Gonzalez CI, Kinzy TG, Dinman JD, Peltz SW: **Mutations in the MOF2/SUI1 gene affect both translation and nonsense-mediated mRNA decay.** *RNA* 1999, **5**:794-804.
68. He F, Jacobson A: **Identification of a novel component of the nonsense-mediated mRNA decay pathway by use of an interacting protein screen.** *Genes Dev* 1995, **9**:437-454.
69. Lykke-Andersen J: **Identification of a human decapping complex associated with hUPF proteins in nonsense-mediated decay.** *Mol Cell Biol* 2002, **22**:8114-8121.
70. Sheth U, Parker R: **Decapping and decay of messenger RNA occur in cytoplasmic processing bodies.** *Science* 2003, **300**:805-808.  
mRNA decay intermediates and degradation proteins are localized to discrete cytoplasmic foci in yeast cells.
71. Bashkurov VI, Scherthan H, Solinger JA, Buerstedde JM, Heyer WD: **A mouse cytoplasmic exoribonuclease with preference for G4 tetraplex substrates.** *J Cell Biol* 1997, **136**:761-773.
72. Ingelfinger D, Ardnt-Jovin DJ, Luhmann R, Achsel T: **The human LSm1-7 proteins co-localize with the mRNA-degrading enzymes Dcp1/2 and Xrn1 in distinct cytoplasmic foci.** *RNA* 2002, **8**:1489-1501.
73. Van Dijk E, Cougot N, Meyer S, Szbajko S, Wahle E, Seraphin B: **Human Dcp2: a catalytically active mRNA decapping enzyme located in specific cytoplasmic structures.** *EMBO J* 2002, **21**:6915-6924.
74. Moore MJ: **RNA events. No end to nonsense.** *Science* 2002, **298**:370-371.